

Synthesis, X-ray Structures, and Reactivity of the First Bis(amino)metallastibanes and Bis(amino)metallabismuthanes

Anne-Marie Caminade*

Laboratoire de Chimie de Coordination du CNRS, Unité No. 8241 liée par conventions à l'Université Paul Sabatier et à l'Institut National Polytechnique, 205 route de Narbonne, F-31077 Toulouse Cedex, France

Michael Veith* and Volker Huch

Institut für Anorganische Chemie der Universität, Im Stadtwald, D-6600 Saarbrücken, West Germany

Wolfgang Malisch*

Institut für Anorganische Chemie der Universität Würzburg, Am Hubland, D-8700 Würzburg, West Germany


Received September 12, 1989

The reaction of $\text{Me}_2\text{Si}(\text{N-}t\text{-Bu})_2\text{ECl}$ ($\text{E} = \text{Sb, Bi}$) with $\text{NaM}(\text{CO})_n\text{Cp}$ ($\text{M} = \text{Fe, Mo, W}$; $\text{Cp} = \eta^5\text{-C}_5\text{H}_5$) affords metallastibanes and bismuthanes, $\text{Cp}(\text{CO})_n\text{ME}(\text{N-}t\text{-Bu})_2\text{SiMe}_2$ (**4a-c**, **5a-c**). The structures of **4c** ($\text{E} = \text{Sb, M} = \text{W}$) and **5c** ($\text{E} = \text{Bi, M} = \text{W}$) have been determined by X-ray crystallography. Both compounds are isotypic and isostructural and crystallize in the triclinic space group with $a = 7.030$ (9) Å, $b = 10.51$ (1) Å, $c = 16.22$ (2) Å, $\alpha = 94.1$ (1)°, $\beta = 90.4$ (1)°, $\gamma = 104.9$ (1)°, $Z = 2$ for **4c** and $a = 7.014$ (9) Å, $b = 10.57$ (1) Å, $c = 16.25$ (2) Å, $\alpha = 94.1$ (1)°, $\beta = 90.1$ (1)°, $\gamma = 104.6$ (1)°, $Z = 2$ for **5c**. The E-W σ -bond lengths are 3.010 (1) Å for $\text{E} = \text{Sb}$ and 3.082 (1) Å for $\text{E} = \text{Bi}$. The reaction of the antimony derivatives with $\text{Fe}_2(\text{CO})_9$ yields $[\text{Fe}(\text{CO})_4]\text{Cp}(\text{CO})_n\text{MSb}(\text{N-}t\text{-Bu})_2\text{SiMe}_2$ (**6a-c**) as a result of the complexation of the antimony lone pair. The crystal structures of **6a** ($\text{M} = \text{Fe}$) and **6b** ($\text{M} = \text{Mo}$) have been determined. **6a** crystallizes in the monoclinic space group with $a = 10.399$ (9) Å, $b = 16.76$ (2) Å, $c = 15.74$ (1) Å, $\beta = 94.16$ (6)°, and $Z = 4$. Both Sb-Fe bond lengths are almost similar: 2.547 (1) Å for the covalent bond and 2.530 (1) Å for the dative bond. **6b** crystallizes in the monoclinic space group with $a = 12.305$ (7) Å, $b = 13.812$ (7) Å, $c = 16.75$ (1) Å, $\beta = 99.03$ (5)°, and $Z = 4$. The Sb-Mo covalent bond length is 2.871 (1) Å and the Sb-Fe dative bond 2.539 (1) Å.

Introduction

Organometallic complexes of the main-group elements are attracting an increasing amount of interest. For example, a significant number of metallaphosphanes, L_nMPR_2 ($\text{M} = \text{transition metal}$) have been synthesized and structurally characterized.¹ However, few investigations deal with the analogous compounds of the heaviest group-15 elements, i.e., metallastibanes² and metallabismuthanes.^{2c,3} Such derivatives have been known since 1964,^{3a} but organic substituents of antimony and bismuth lack diversification as in most cases, alkyl or phenyl groups, sometimes halides, but no amino groups have been used. Moreover, only one structure, $\text{PPh}_3(\text{CO})_3\text{CoBiPh}_2$, has

Scheme I



been determined by X-ray diffraction so far.^{3e}

This paper reports the synthesis of the first amino-substituted metallastibanes and bismuthanes, their single-crystal X-ray diffraction characterization, and their behavior toward $\text{Fe}_2(\text{CO})_9$.

Results and Discussion

Synthesis of Metallastibanes and Metallabismuthanes. The reaction of $\text{Me}_2\text{Si}(\text{N-}t\text{-Bu})_2\text{ECl}$, **1** ($\text{E} = \text{Sb}$) and **2** ($\text{E} = \text{Bi}$),⁴ in hexane or benzene solution with $\text{NaM}(\text{CO})_n\text{Cp}$, **3a-c** ($\text{M} = \text{Fe, Mo, W}$), at room temperature yields, by a nucleophilic metalation process, the corresponding cyclic bis(amino)metallastibanes, **4a-c**, and -bismuthanes, **5a-c** (Scheme I).

¹H NMR spectroscopy shows that crude **4a-c** and **5a-c** are quantitatively obtained. Compounds **4a-c** and **5b-c** are isolated in 46–71% yield as extremely air- and moisture-sensitive powders or crystals. This sensitivity is related at least in part to the high lability of the E-N bonds, which are easily cleaved to give bis(*tert*-butylamino)dimethylsilane. The most labile compound is **5a**, which has not been isolated but only characterized in solution. Its ¹H NMR spectrum exhibits four singlets in the range expected for such a compound:⁴ 0.19 and 0.38 ppm for the

(1) For X-ray structures of R_2PM compounds, see, for example: (a) Barrow, M. J.; Sim, G. A.; Dobbie, R. C.; Mason, P. R. *J. Organomet. Chem.* 1974, 69, C4. (b) Barrow, M. J.; Sim, G. A. *J. Chem. Soc., Dalton Trans.* 1975, 291. (c) Hutchins, L. D.; Duesler, E. N.; Paine, R. T. *Organometallics* 1982, 1, 1254. (d) Baker, R. T.; Whitney, J. F.; Wreford, S. S. *Organometallics* 1983, 2, 1049. (e) Ebsworth, E. A. V.; Gould, R. O.; Mc Manus, N. T.; Pilkington, N. J.; Rankin, D. W. H. *J. Chem. Soc., Dalton Trans.* 1984, 2561. (f) Bohle, D. S.; Jones, T. C.; Rickard, C. E. F.; Roper, W. R. *J. Chem. Soc., Chem. Commun.* 1984, 865. (g) Malisch, W.; Angerer, W.; Cowley, A. H.; Norman, N. C. *J. Chem. Soc., Chem. Commun.* 1985, 1811. (h) Arif, A. M.; Cowley, A. H.; Nunn, C. M.; Pakulski, M. *J. Chem. Soc., Chem. Commun.* 1987, 994.

(2) (a) Cullen, W. R.; Patmore, D. J.; Sams, J. R.; Newlands, M. J.; Thompson, L. K. *J. Chem. Soc., Chem. Commun.* 1971, 952. (b) Malisch, W.; Panster, P. *J. Organomet. Chem.* 1974, 76, C7. (c) Malisch, W.; Panster, P. *J. Organomet. Chem.* 1975, 99, 421, and references therein. (d) Kaul, H. A.; Greissing, D.; Luksza, M.; Malisch, W. *J. Organomet. Chem.* 1982, 228, C29. (e) Thum, G.; Ries, W.; Greissing, D.; Malisch, W. *J. Organomet. Chem.* 1983, 252, C67. (f) Wade, S. R.; Wallbridge, M. G. H.; Willey, G. P. *J. Organomet. Chem.* 1984, 267, 271. (g) Schemm, R.; Malisch, W. *J. Organomet. Chem.* 1985, 288, C9.

(3) (a) Nesmeyanov, A. N.; Anisimov, K. N.; Kolobova, N. E.; Khandozhko, V. N. *Dokl. Akad. Nauk. SSSR* 1964, 156, 383 (CA61; 7041d). (b) Shrauzer, G. N.; Kratel, G. *Angew. Chem.* 1965, 77, 130; *Angew. Chem., Int. Ed. Engl.* 1965, 4, 146. (c) Schrauzer, G. N.; Kratel, G. *Chem. Ber.* 1969, 102, 2392. (d) Cullen, W. R.; Patmore, D. J.; Sams, J. R. *Inorg. Chem.* 1973, 12, 867. (e) Panster, P.; Malisch, W. *J. Organomet. Chem.* 1977, 134, C32. (f) Calderazzo, F.; Poli, R.; Pellizzi, G. *J. Chem. Soc., Dalton Trans.* 1984, 2535.

(4) Veith, M.; Bertsch, B. Z. *Anorg. Allg. Chem.* 1988, 557, 7.

Table I. $^1\text{H NMR}^a$ and IR b Data for Compounds 4a-c, 5a-c, and 6a-c

compound	$^1\text{H NMR}, \delta$			IR, $\nu(\text{CO})$
	Me	tBu	Cp	
$\text{Me}_2\text{Si}(\text{N}-t\text{-Bu})_2\text{SbFe}(\text{CO})_2\text{Cp}$ 4a	0.35 0.47	1.19	4.28	1982 (s), 1933 (s)
$\text{Me}_2\text{Si}(\text{N}-t\text{-Bu})_2\text{SbMo}(\text{CO})_3\text{Cp}$ 4b	0.46	1.20	4.73	1985 (s), 1917 (m), 1888 (s)
$\text{Me}_2\text{Si}(\text{N}-t\text{-Bu})_2\text{SbW}(\text{CO})_3\text{Cp}$ 4c	0.41 0.45	1.19	4.77	1987 (s), 1916 (m), 1883 (s)
$\text{Me}_2\text{Si}(\text{N}-t\text{-Bu})_2\text{BiFe}(\text{CO})_2\text{Cp}$ 5a	0.19 0.38	1.13	4.45	1976 (s), 1929 (s)
$\text{Me}_2\text{Si}(\text{N}-t\text{-Bu})_2\text{BiMo}(\text{CO})_3\text{Cp}$ 5b	0.24 0.35	1.14	4.86	1982 (s), 1910 (w), 1878 (s)
$\text{Me}_2\text{Si}(\text{N}-t\text{-Bu})_2\text{BiW}(\text{CO})_3\text{Cp}$ 5c	0.21 0.35	1.14	4.95	1983 (s), 1909 (w), 1878 (m)
$\text{Me}_2\text{Si}(\text{N}-t\text{-Bu})_2\text{Sb}$ 6a	$\text{Fe}(\text{CO})_2\text{Cp}$			2039 (s), 2012 (s), 1977 (br), 1954 (w), 1924 (br s)
	$\text{Fe}(\text{CO})_4$	0.33		
$\text{Me}_2\text{Si}(\text{N}-t\text{-Bu})_2\text{Sb}$ 6b	$\text{Mo}(\text{CO})_3\text{Cp}$			2038 (m), 2012 (s), 1947 (br s), 1927 (m), 1911 (w)
	$\text{Fe}(\text{CO})_4$	0.42	1.31 4.74	
$\text{Me}_2\text{Si}(\text{N}-t\text{-Bu})_2\text{Sb}$ 6c	$\text{W}(\text{CO})_3\text{Cp}$	0.42		2038 (m), 2009 (s), 1939 (br s), 1921 (m), 1906 (w)
	$\text{Fe}(\text{CO})_4$	0.59	1.30 4.75	

^aSpectra were recorded on a solution of the compound in C_6D_6 .
^bHexane solution for 4a-c and 5a-c, benzene for 6a-c.




Figure 1. ORTEP drawing of 4c. Important bond distances (Å) and angles (deg): SbW = 3.010 (1); N(1)Sb = 2.050 (7); N(2)Sb = 2.066 (7); N(1)Si = 1.734 (6); N(2)Si = 1.733 (8); N(1)SbW = 105.7 (2); N(2)SbW = 105.0 (2); N(1)SbN(2) = 74.3 (3); N(1)SiN(2) = 91.6 (3); SiN(1)Sb = 96.9 (3); SiN(2)Sb = 96.4 (3).

two magnetically nonequivalent methyl groups, 1.13 ppm for the two equivalent *tert*-butyl groups, and 4.45 ppm for the η^5 -bonded cyclopentadienyl ring. Furthermore, two intense carbonyl stretching vibrations are present in the IR spectrum of 5a as in the spectrum of the analogous isolated antimony derivative, 4a. Similarly, the IR spectra of 4b,c and 5b,c exhibit three carbonyl stretching vibrations typical of a C_s symmetry ($A'(1)$, $A'(2)$, and A'' ; Table I).

To ascertain further the structure of the compounds of types 4 and 5, X-ray structure determinations have been undertaken for 4c and 5c.

X-ray Structural Analysis of 4c and 5c. The results of these determinations appear in Figures 1 for 4c and 2 for 5c, together with the atom-numbering protocol and selected bond lengths and angles. Pertinent crystallographic data are collected in Table II (compounds 4c and




Figure 2. ORTEP drawing of 5c. Important bond distances (Å) and angles (deg): BiW = 3.082 (1); N(1)Bi = 2.17 (1); N(2)Bi = 2.18 (1); N(1)Si = 1.75 (1); N(2)Si = 1.74 (1); N(1)BiW = 104.4 (3); N(2)BiW = 103.1 (4); N(1)BiN(2) = 71.4 (4); N(1)SiN(2) = 93.5 (6); SiN(1)Bi = 96.9 (5); SiN(2)Bi = 96.9 (5).

Table II. Summary of Crystal Data, Intensity Collection and Structure Solution and Refinement for 4c and 5c

	4c	5c
formula	$\text{C}_{18}\text{H}_{29}\text{N}_2\text{O}_3\text{SbSiW}$	$\text{C}_{18}\text{H}_{29}\text{BiN}_2\text{O}_3\text{SiW}$
space group	$P\bar{1}$	$P\bar{1}$
cryst syst	triclinic	triclinic
a , Å	7.030 (9)	7.014 (9)
b , Å	10.51 (1)	10.57 (1)
c , Å	16.22 (2)	16.25 (2)
α , deg	94.1 (1)	94.1 (1)
β , deg	90.4 (1)	90.1 (1)
γ , deg	104.9 (1)	104.6 (1)
Z	2	2
V , Å ³	1155	1164
d_{calc} , g/cm ³	1.778	2.012
cryst dims, mm ³	0.20 × 0.15 × 0.30	0.25 × 0.25 × 0.35
radiation	$\text{Mo K}\alpha$ (0.710 69 Å)	$\text{Mo K}\alpha$ (0.710 69 Å)
μ , cm ⁻¹	63.06	126.50
no. of unique reflns measd	2958	2750
no. of obsd reflns for refinement	2769 ($F_o > 3\sigma(F_o)$)	2136 ($F_o > 3\sigma(F_o)$)
no. of variables used in refinement	221	220
R	0.050	0.044
R_w	0.052	0.045
$w = k/(\delta^2_F + g F^2)$	$k, 1.0; g, 0.010 13$	$k, 0.6864; g, 0.005 65$
atoms refined anisotropically	W, Sb, Si, O, N, C1, C5, C11-C13	W, Bi, Si, O, N, C1, C5, C11-C13

5c are isotypic and isostructural).

Both crystal structures consist of distinct, well-separated monomers, which are in perfect agreement with our formulation. The four-membered rings N(1)SiN(2)E (E = Sb or Bi) are not as perfectly planar as in 1 and 2⁴ (dihedral angles between N(1)SiN(2) and N(1)EN(2) planes (E = Sb, Bi) are in 4c 9.7° and in 5c 10.4°) and the nitrogen atoms are slightly pyramidal: the sum of angles around N(1) and N(2) is 358.4° and 352.3° for 4c, 357.9° and 350.9° for 5c. The two *tert*-butyl groups are in the *cis* position relative to the plane of the cycle, due to steric interactions between hydrogen atoms of the cyclopentadienyl ring and the *tert*-butyl groups (4c, C(2)···C(16), C(2)···C(17) = 3.8 Å; 5c, C(2)···C(16) = 3.8 Å, C(2)···C(17) = 4.0 Å. This indicates that the methyl group occupies the gap between C(16) and C(17)). We believe that these steric repulsions are also responsible for the larger sum of the angles found at antimony (285°) or bismuth (278.9°)

Scheme II





Figure 3. ORTEP drawing of **6b**. Important bond distances (Å) and angles (deg): MoSb = 2.871 (1); FeSb = 2.539 (1); N(1)Sb = 2.054 (7); N(2)Sb = 2.050 (6); N(1)Si = 1.736 (7); N(2)Si = 1.725 (8); MoSbFe = 122.8 (1); N(1)SbMo = 108.9 (2); N(2)SbMo = 111.5 (2); N(1)SbFe = 115.0 (2); N(2)SbFe = 114.1 (2); N(1)SbN(2) = 74.5 (3); N(1)SiN(2) = 91.7 (3); SiN(1)Sb = 96.6 (3); SiN(2)Sb = 97.1 (3).

with respect to 1 (274.5°) or 2 (267.4°).

To the best of our knowledge, **4c** represents the first structurally characterized example of stibido ligands (and **5c** the second for bismuthido ligands) terminally coordinated to transition-metal centers, and therefore direct comparisons of E–W distances are not possible. The observed E–W distances (3.010 Å for E = Sb and 3.082 Å for E = Bi), although much longer than those described in the literature for dative σ -bonds (e.g., 2.757 and 2.766 Å for $[\text{W}(\text{CO})_5]_2[\mu\text{-W}(\text{CO})_5](\text{SbPh})_2$ ⁵ and 2.851 Å for $[\text{W}(\text{CO})_5\text{BiMe}][\text{W}_2(\text{CO})_8\text{Bi}_2]$ ⁶, are close to the sum of covalent radii of antimony or bismuth (1.40 and 1.46 Å, respectively) and of tungsten (1.611 Å) obtained from Cp(CO)₃W–W(CO)₃Cp.⁷ In comparison to the chlorine compounds **1** and **2**,⁴ the Si–N distances in **4c** and **5c** are of the same magnitude (1.73–1.75 Å).

Complexation of Metallastibanes. When a benzene solution of metallastibanes **4a–c** is added to a benzene suspension of Fe₂(CO)₉, the corresponding complexes **6a–c** are isolated in 59–62% yield as relatively air stable compounds (Scheme II).

The reaction of metallabismuthanes **5a–c** under the same experimental conditions did not afford the corresponding metallabismuthane complexes. This illustrates the well-known decrease in the availability for bonding of




Figure 4. ORTEP drawing of **6a**. Important bond distances (Å) and angles (deg): Fe(1)Sb = 2.547 (1); Fe(2)Sb = 2.530 (1); N(2)Sb = 2.042 (5); N(1)Sb = 2.045 (5); N(2)Si = 1.736 (6); N(1)Si = 1.735 (6); Fe(1)SbFe(2) = 121.1 (1); N(2)SbFe(1) = 110.2 (2); N(1)SbFe(1) = 110.4 (2); N(2)SbFe(2) = 115.5 (2); N(1)SbFe(2) = 115.9 (2); N(2)SbN(1) = 74.7 (2); N(2)SiN(1) = 91.2 (3); SiN(2)Sb = 97.0 (3); SiN(1)Sb = 96.9 (3).

Table III. Summary of Crystal Data, Intensity Collection, and Structure Solution and Refinement for **6a** and **6b**

	6a	6b
formula	C ₂₁ H ₂₉ Fe ₂ N ₂ O ₆ SbSi	C ₂₂ H ₂₉ FeMoN ₂ O ₇ SbSi
space group	P2 ₁ /n	Cc
cryst syst	monoclinic	monoclinic
a, Å	10.399 (9)	12.305 (7)
b, Å	16.76 (2)	13.812 (7)
c, Å	15.74 (1)	16.75 (1)
β, deg	94.16 (6)	99.03 (5)
Z	4	
V, Å ³	2736	2812
d _{calc} , g/cm ³	1.619	1.736
cryst dimens, mm ³	0.10 × 0.12 × 0.20	0.12 × 0.15 × 0.25
radiation	M _o Kα (0.71069 Å)	M _o Kα (0.71069 Å)
μ, cm ⁻¹	20.84	19.52
no. of unique reflns measd	3374	2188
no. of obsd reflns for refinement	2923 (F _o > 2σ(F _o))	2020 (F _o > 2σ(F _o))
variables used in refinement	323	269
R	0.033	0.027
R _w		0.033
w = k/(σ ² F + g F ²)	k, unit weight; g, unit weight	k, 1.0; g, 0.00135
atoms refined anisotropically	Sb, Fe, Si, O, N, C1–C21	Sb, Mo, Fe, Si, O, N, C1–C7, C13, C17

the lone pair with the group-15 elements, on going from phosphorus to bismuth, and may be related to an increase in the s character of the lone pair.

Compounds **6a–c** have been fully characterized. A comparison of ¹H NMR values (Table I) of free, **4**, and complexed, **6**, derivatives shows, as expected,^{2c} a downfield shift of the methyl and *tert*-butyl resonances as a result of complexation. A downfield shift is also observed for the cyclopentadienyl signal of **6a**, whereas the corresponding value for **6b,c** remains essentially unchanged.

X-ray diffraction studies have been undertaken to confirm the complexation of the lone pair at antimony.

X-ray Structural Analysis of 6a and 6b. The structures of **6b** and **6a** are illustrated in Figures 3 and 4, respectively, together with atom-numbering schemes and


(5) Huttner, G.; Weber, U.; Sigwarth, B.; Scheidsteger, O. *Angew. Chem.* **1982**, *94*, 210; *Angew. Chem., Int. Ed. Engl.* **1982**, *21*, 215; *Angew. Chem. Suppl.* **1982**, 411.

(6) Arif, A. M.; Cowley, A. H.; Norman, N. C.; Pakulski, M. *J. Am. Chem. Soc.* **1985**, *107*, 1062.

(7) Adams, R. D.; Collins, D. M.; Cotton, F. A. *Inorg. Chem.* **1974**, *13*, 1086.

selected bond lengths and angles. Pertinent crystallographic data are summarized in Table III.

Figures 3 and 4 support our assumptions concerning the complexation of the antimony lone pair by $\text{Fe}(\text{CO})_4$. The environment around the iron (tetracarbonyl) atom is a distorted trigonal bipyramid in both structures. The largest deviation from ideality is found for the $\text{SbFeC}_{\text{axial}}$ angles: 174.5° for **6a** and 175.9° for **6b**. Among the six angles around antimony, only two are far from tetrahedral values: the FeSbM angles (121.1° for **6a** ($M = \text{Fe}$), 122.8° for **6b** ($M = \text{Mo}$)) which are larger than the value found in the related compound **7** ($\text{FeSbFe} = 117^\circ$)⁸ and the



$\text{N}(1)\text{SbN}(2)$ angles which are identical for the free and complexed derivatives (74.3° for **4c**, 74.7° for **6a**, 74.5° for **6b**). On the other hand, complexation affects the average NSbM angles: 105.3° for uncomplexed **4c**, 110.3° and 110.2° for **6a** and **6b**, respectively.

Another interesting feature of the structure of **6a,b** is the covalent bond length $\text{Sb}-\text{M}$. If the length is calculated as for **4c** (i.e., the sum of the molybdenum covalent radius (1.617 Å) obtained from $\text{Cp}_2\text{Mo}_2(\text{CO})_6$ ⁷ and of the antimony covalent radius (1.40 Å)), the result is 3.017 Å, whereas the actual bond length is 2.871 Å for **6b**. Thus, complexation of the antimony lone pair induces a dramatic shortening of the $\text{Sb}-\text{M}$ covalent bond length. This is true also for **6a** since both $\text{Sb}-\text{Fe}$ bond lengths (2.547 Å for the covalent bond and 2.530 Å for the dative bond) are almost equal. We have interpreted the shortening of the covalent $\text{Sb}-\text{M}$ bond in terms of an increased amount of $\text{M}\rightarrow\text{Sb}$ $d\pi-d\pi$ back-bonding (the similarity of the $\text{Sb}-\text{N}$ bond lengths in **4c** and **6a,b** argues against any noticeable decrease in the σ -bonding covalent radius of antimony). This is also consistent with the downfield shift of the cyclopentadienyl resonance observed in the ^1H NMR spectrum of **6a** in comparison to **4a**.

Experimental Section

All operations were carried out under an atmosphere of dried nitrogen. Solvents were refluxed over lithium aluminum hydride for 3 days before distillation and kept over sodium wires. $\text{Me}_2\text{Si}(\text{N}-t\text{-Bu})_2\text{SbCl}$, **1**, and $\text{Me}_2\text{Si}(\text{N}-t\text{-Bu})_2\text{BiCl}$, **2**, were prepared as described in the literature,⁴ as was **3a-c**.⁹ ^1H NMR spectra were recorded on a Bruker WP-80 instrument. ^1H chemical shifts are reported in parts per million relative to Me_2Si . Infrared spectra were recorded on a Perkin-Elmer 883 using one NaCl cell, the solvent spectrum being subtracted from the complex solution spectrum. Melting points were measured in sealed capillaries. Molecular weight measurements were carried out in benzene.

General Procedure for the Synthesis of $[\text{Me}_2\text{Si}(\text{N}-t\text{-Bu})_2]\text{EM}(\text{CO})_n(\eta^5\text{-C}_5\text{H}_5)$, **4a-c and **5a-c**.** Typically a hexane solution of $[\text{Me}_2\text{Si}(\text{N}-t\text{-Bu})_2]\text{ECl}$ ($\sim 10^{-3}$ mol in 30 mL) was added dropwise at room temperature to a suspension of $\text{NaM}(\text{CO})_n\text{Cp}$ ($\sim 2 \times 10^{-3}$ mole, 70–100% excess) in hexane (10 mL). The resulting mixture was magnetically stirred overnight. NaCl and excess $\text{NaM}(\text{CO})_n\text{Cp}$ then were removed by filtration. An extremely air- and moisture-sensitive red solution of $\text{Cp}(\text{CO})_n\text{ME}(\text{N}-t\text{-Bu})_2\text{SiMe}_2$, **4a-c** or **5a-c**, was obtained. According to ^1H NMR spectra, yields were nearly quantitative.

Synthesis of 1,3-Di-tert-butyl-2,2-dimethyl-4-[(dicarbonyl- η^5 -cyclopentadienyl)ferrio]-1,3,2,4-diazasilastibetidine, $\text{Cp}(\text{CO})_2\text{FeSb}(\text{N}-t\text{-Bu})_2\text{SiMe}_2$, **4a.** The solution of **4a** obtained from 0.291 g (0.815 mmol) of $\text{Me}_2\text{Si}(\text{N}-t\text{-Bu})_2\text{SbCl}$, **1**,

and 0.277 g (1.38 mmol) of $\text{NaFe}(\text{CO})_2\text{Cp}$, **3a**, was concentrated under vacuum to ~ 2 mL and cooled for 1 h at -78°C . **4a** precipitated as an orange powder (0.240 g, 0.481 mmol, 59% yield), mp $103\text{--}105^\circ\text{C}$. Calcd mol wt 499 g/mol, found (cryoscopic) 501 g/mol. Anal. Calcd for $\text{C}_{17}\text{H}_{25}\text{FeN}_2\text{O}_2\text{SbSi}$: C, 40.90; H, 5.87; N, 5.61. Found: C, 42.49; H, 6.26; N, 4.88.

Synthesis of 1,3-Di-tert-butyl-2,2-dimethyl-4-[(tricarbonyl- η^5 -cyclopentadienyl)molybdenio]-1,3,2,4-diazasilastibetidine, $\text{Cp}(\text{CO})_3\text{MoSb}(\text{N}-t\text{-Bu})_2\text{SiMe}_2$, **4b.** The solution of **4b** obtained from 0.397 g (1.11 mmol) of $\text{Me}_2\text{Si}(\text{N}-t\text{-Bu})_2\text{SbCl}$, **1**, and 0.589 g (2.20 mmol) of $\text{NaMo}(\text{CO})_3\text{Cp}$, **3b**, was concentrated under vacuum to ~ 2 mL and left for 1 day at -20°C . **4b** crystallized as small brown needles (0.289 g, 0.511 mmol, 46% yield), mp $78\text{--}80^\circ\text{C}$. Calcd mol wt 567 g/mol, found (cryoscopic) 569 g/mol. Anal. Calcd for $\text{C}_{18}\text{H}_{26}\text{MoN}_2\text{O}_3\text{SbSi}$: C, 38.11; H, 5.16; N, 4.94. Found: C, 38.13; H, 5.33; N, 4.95.

Synthesis of 1,3-Di-tert-butyl-2,2-dimethyl-4-[(tricarbonyl- η^5 -cyclopentadienyl)tungstenio]-1,3,2,4-diazasilastibetidine, $\text{Cp}(\text{CO})_3\text{WSb}(\text{N}-t\text{-Bu})_2\text{SiMe}_2$, **4c.** A 0.520-g (1.46 mmol) sample of $\text{Me}_2\text{Si}(\text{N}-t\text{-Bu})_2\text{SbCl}$, **1**, and 0.964 g (2.71 mmol) of $\text{NaW}(\text{CO})_3\text{Cp}$, **3c**, were reacted to give a solution of **4c**, which was concentrated to ~ 15 mL and left for 4 h at room temperature. **4c** was obtained as orange crystals (0.681 g, 1.04 mmol, 71% yield), mp $139\text{--}141^\circ\text{C}$. Anal. Calcd for $\text{C}_{18}\text{H}_{26}\text{W}_2\text{N}_2\text{O}_3\text{SbSi}$: C, 32.99; H, 4.47; N, 4.28. Found: C, 32.92; H, 4.36; N, 4.25. One of these crystals was suitable for an X-ray structure determination.

Synthesis of 1,3-Di-tert-butyl-2,2-dimethyl-4-[(dicarbonyl- η^5 -cyclopentadienyl)ferrio]-1,3,2,4-diazasilabismuthetidine, $\text{Cp}(\text{CO})_2\text{FeBi}(\text{N}-t\text{-Bu})_2\text{SiMe}_2$, **5a.** The reaction was carried out with 0.237 g (0.533 mmol) of $\text{Me}_2\text{Si}(\text{N}-t\text{-Bu})_2\text{BiCl}$, **2**, and 0.203 g (1.01 mmol) of $\text{NaFe}(\text{CO})_2\text{Cp}$, **3a**, but all attempts to isolate **5a** failed. This compound was characterized only by ^1H NMR and IR spectra (Table I).

Synthesis of 1,3-Di-tert-butyl-2,2-dimethyl-4-[(tricarbonyl- η^5 -cyclopentadienyl)molybdenio]-1,3,2,4-diazasilabismuthetidine, $\text{Cp}(\text{CO})_3\text{MoBi}(\text{N}-t\text{-Bu})_2\text{SiMe}_2$, **5b.** The solution of **5b** obtained from 0.38 g (0.855 mmol) of $\text{Me}_2\text{Si}(\text{N}-t\text{-Bu})_2\text{BiCl}$, **2**, and 0.441 g (1.65 mmol) of $\text{NaMo}(\text{CO})_3\text{Cp}$, **3b**, was slowly concentrated under vacuum. Small brown needles of **5b** were deposited on the walls of the flask (0.308 g, 0.47 mmol, 55% yield), mp $130\text{--}132^\circ\text{C}$. Calcd mol wt 645.5 g/mol, found (cryoscopic) 682 g/mol (probably because of decomposition). Anal. Calcd for $\text{C}_{18}\text{H}_{26}\text{BiMoN}_2\text{O}_3\text{Si}$: C, 33.08; H, 4.48; N, 4.28. Found: C, 33.14; H, 4.60; N, 4.12.

Synthesis of 1,3-Di-tert-butyl-2,2-dimethyl-4-[(tricarbonyl- η^5 -cyclopentadienyl)tungstenio]-1,3,2,4-diazasilabismuthetidine, $\text{Cp}(\text{CO})_3\text{WBi}(\text{N}-t\text{-Bu})_2\text{SiMe}_2$, **5c.** $\text{Me}_2\text{Si}(\text{N}-t\text{-Bu})_2\text{BiCl}$, **2** (0.418 g, 0.941 mmol), and 0.655 g (1.84 mmol) of $\text{NaW}(\text{CO})_3\text{Cp}$, **3c**, were reacted to give a solution of **5c**, which was concentrated to ~ 10 mL and left for 3 days at -20°C . **5c** was obtained as red crystals (0.363 g, 0.489 mmol, 52% yield). This compound decomposed without melting at $\sim 150^\circ\text{C}$ (the color changed from red to black). Anal. Calcd for $\text{C}_{18}\text{H}_{26}\text{BiN}_2\text{O}_3\text{SiW}$: C, 29.12; H, 3.95; N, 3.77. Found: C, 29.28; H, 4.06; N, 3.67.

A single crystal was grown by cooling slowly (32 h) a solution of **5c** (~ 0.1 mmol) in hexane (2 mL) from room temperature to -20°C .

General Procedure for the Synthesis of $[\text{Fe}(\text{CO})_4]\text{Cp}(\text{CO})_n\text{MSb}(\text{N}-t\text{-Bu})_2\text{SiMe}_2$, **6a-c.** An hexane solution of **4a-c** (~ 0.5 mmol) obtained as described above was evaporated to dryness. The resulting powder was solved in benzene (20 mL) and used without further purification. This benzene solution was added dropwise to a suspension of $\text{Fe}_2(\text{CO})_9$ (~ 0.5 mmol) in benzene (10 mL), and the resulting mixture was stirred overnight. The dark-red solution thus obtained was filtered to remove small amounts of insoluble materials and then evaporated to dryness. A relatively air-stable brown powder of **6a-c** was obtained.

An identical procedure applied to $\text{Cp}(\text{CO})_3\text{WBi}(\text{N}-t\text{-Bu})_2\text{SiMe}_2$, **5c**, did not afford the corresponding complex. **5c** and $\text{Fe}_2(\text{CO})_9$ were recovered unreacted.

Synthesis of Tetracarbonyl[1,3-di-tert-butyl-2,2-dimethyl-4-[(dicarbonyl- η^5 -cyclopentadienyl)ferrio]-1,3,2,4-diazasilastibetidine]iron, $[\text{Fe}(\text{CO})_4]\text{Cp}(\text{CO})_2\text{FeSb}(\text{N}-t\text{-Bu})_2\text{SiMe}_2$, **6a.** The powder of **6a** obtained from 0.171 g (0.479 mmol) of $\text{Me}_2\text{Si}(\text{N}-t\text{-Bu})_2\text{SbCl}$, **1**, 0.184 g (0.921 mmol) of

(8) Cane, D. J.; Forbes, E. J.; Hamor, T. A. *J. Organomet. Chem.* 1976, 117, C101.

(9) Wilkinson, G.; Piper, T. S. *J. Inorg. Nucl. Chem.* 1956, 3, 104.

NaFe(CO)₂Cp **3a**, and 0.171 g (0.471 mmol) of Fe₂(CO)₉ was dissolved in a minimum amount of toluene at room temperature; then the solution was cooled at -20 °C, affording red crystals of **6a** (0.198 g, 0.297 mmol, 62% yield based upon 1), mp 238-240 °C. Anal. Calcd for C₂₁H₂₉Fe₂N₂O₆SbSi: C, 37.81; H, 4.39; N, 4.20. Found: C, 38.21; H, 4.53; N, 4.25.

Some crystals (~10⁻⁵ mol) were dissolved in refluxing hexane (25 mL); the solution was filtered, cooled to 0 °C, and left overnight at this temperature, affording red single crystals.

Synthesis of Tetracarbonyl[1,3-di-*tert*-butyl-2,2-dimethyl-4-[(tricarboxyl-η⁵-cyclopentadienyl)molybdenio]-1,3,2,4-diazasilastibetidine]iron, [Fe(CO)₄]Cp(CO)₃MoSb(N-*t*-Bu)₂SiMe₂, **6b.** Hexane (5 mL) was added to the powder of **6b** obtained from 0.208 g (0.582 mmol) of Me₂Si(N-*t*-Bu)₂SbCl, 1, 0.292 g (1.09 mmol) of NaMo(CO)₃Cp, **3b**, and 0.209 g (0.575 mmol) of Fe₂(CO)₉. The resulting brown suspension was vigorously magnetically stirred and then allowed to settle. Solvent was removed by aid of a pipet, and hot toluene (3 mL) was added to the insoluble material. The resulting mixture was filtered to give a red solution, which was cooled overnight at -20 °C. **6b** was obtained as red crystals (0.269 g, 0.366 mmol, 63% yield based upon 1), mp 230-232 °C. Anal. Calcd for C₂₂H₂₉FeMoN₂O₇SbSi: C, 35.94; H, 3.98; N, 3.81. Found: C, 36.47; H, 4.06; N, 3.74.

Crystals of **6b** were dissolved in a minimum amount of hot toluene, and the resulting solution was slowly cooled overnight from 100 °C to room temperature, affording single crystals of **6b**.

Synthesis of Tetracarbonyl[1,3-di-*tert*-butyl-2,2-dimethyl-4-[(tricarboxyl-η⁵-cyclopentadienyl)tungstenio]-1,3,2,4-diazasilastibetidine]iron, [Fe(CO)₄]Cp(CO)₃WSb(N-*t*-Bu)₂SiMe₂, **6c.** The brown powder of **6c** obtained from 0.201 g (0.563 mmol) of Me₂Si(N-*t*-Bu)₂SbCl, 1, 0.391 g (1.10 mmol) of NaW(CO)₃Cp, **3c**, and 0.201 g (0.552 mmol) of Fe₂(CO)₉ was dissolved in toluene (~15 mL) and cooled at -20 °C. **6c** was obtained as a red powder (0.273 g, 0.332 mmol, 59% yield based upon 1). **6c** is not soluble enough in benzene to allow molecular weight measurement; mp 227-229 °C. Anal. Calcd for C₂₂H₂₉FeN₂O₇SbSiW: C, 32.10; H, 3.56; N, 3.40. Found: C, 32.62; H, 3.65; N, 3.42.

X-ray Studies. Crystal Structure Determinations. Suitable crystals of each compound **4c**, **5b**, **6a**, and **6b** were examined by using similar procedures.

The crystallographic data are summarized in Tables II and III. Unit cell parameters were determined from 18-26 carefully centered reflections in the range 20 < 2θ < 25°. Data were collected by using the ω-θ scan technique up to a 45° 2θ angle. An empirical absorption correction based on a series of ψ scans was made.

The structures were solved by Patterson and difference Fourier techniques and refined by full-matrix least-squares analysis with anisotropic thermal parameters for all metal, oxygen, nitrogen, and some carbon atoms (see Tables II and III).

All hydrogen atoms were fixed at their carbon atoms and refined together as rigid groups. In **6b** the Cp ring (C(8)-C(12)) was refined as a fixed regular pentagon. One of the *tert*-butyl groups (C18-C20) is disordered, so each carbon atom occupies two positions with a half weight.

All calculations were carried out on a DEC Micro VAX II computer using the SHELXS-86 and SHELX-76 programs.¹¹

Acknowledgment. We thank the Alexander von Humboldt Foundation for a Postdoctoral fellowship (A.M.C.).

Registry No. 1, 118083-14-0; 2, 118083-15-1; **3a**, 12152-20-4; **3b**, 12107-35-6; **3c**, 12107-36-7; **4a**, 126457-14-5; **4b**, 126457-15-6; **4c**, 126457-16-7; **5a**, 126422-15-9; **5b**, 126422-16-0; **5c**, 126422-17-1; **6a**, 126457-17-8; **6b**, 126457-18-9; **6c**, 126457-19-0.

Supplementary Material Available: Tables of atomic coordinates, bond distances and angles, and the anisotropic temperature values of the atoms of structures **4c**, **5c**, **6a**, and **6b** (12 pages); listings of structure factor amplitudes of **4c**, **5c**, **6a**, and **6b** (28 pages). Ordering information is given on any current masthead page.

(10) Hamilton, W. C. *Acta Crystallogr.* 1959, 12, 609.

(11) Sheldrick, G. Program for Crystal Structure Determination, Version SHELX-76 and SHELXS-86.

Ligand-Field and Charge-Transfer Photochemistry of M(CO)₄(1,10-phenanthroline) (M = Cr, Mo, W). Mechanistic Information from High-Pressure Effects

S. Wieland,¹ K. Bal Reddy,² and R. van Eldik*

Institute for Inorganic Chemistry, University of Witten/Herdecke, Stockumer Strasse 10, 5810 Witten, FRG

Received October 12, 1989

The pressure dependence of the quantum yield for photosubstitution of CO in M(CO)₄(1,10-phenanthroline) (M = Cr, Mo, W) was studied at different excitation wavelengths in order to resolve the nature of the substitution mechanism operating from the ligand-field and charge-transfer excited states. Irradiation at 366 nm results in positive volumes of activation (between +6 and +10 cm³ mol⁻¹) for ligand-field photosubstitution, in agreement with a dissociative substitution mode. Irradiation at 546 nm results in significantly negative volumes of activation (between -12 and -14 cm³ mol⁻¹) for charge-transfer photosubstitution in the case of M = Mo and W, which is interpreted as strong evidence for an associative substitution reaction originating from the MLCT excited state. The corresponding value is small positive for the Cr complex and ascribed to a dissociative process operating from the charge-transfer excited state.

Introduction

The application of high-pressure kinetic techniques in mechanistic studies on the reactions of inorganic and organometallic complexes in solution has received much

attention from various groups in recent years.³⁻⁶ Especially the application of this method in the elucidation of the mechanisms of thermal reactions has been rather successful. We and others⁷ have in recent years investi-

(1) Work performed at the Institute for Physical and Theoretical Chemistry, University of Frankfurt, FRG.

(2) On leave from the Department of Chemistry Osmania University, Hyderabad-500007, India.

(3) van Eldik, R., Ed. *Inorganic High Pressure Chemistry: Kinetics and Mechanisms*; Elsevier: Amsterdam, 1986.

(4) Merbach, A. E. *Pure Appl. Chem.* 1987, 59, 161.

(5) Kotowski, M.; van Eldik, R. *Coord. Chem. Rev.* 1989, 93, 19.

(6) van Eldik, R.; Asano, T.; le Noble, W. J. *Chem. Rev.* 1989, 89, 549.